

University of Groningen

A genomic perspective timely needed for re-evaluating the species delimitations, evolutionary trajectories, and conservation strategies of the Galapagos giant tortoises

Fontaine, Michael

Published in:
Peer Community in Evolutionary Biology

DOI:
[10.24072/pci.evolbiol.100031](https://doi.org/10.24072/pci.evolbiol.100031)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2017

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Fontaine, M. (2017). A genomic perspective timely needed for re-evaluating the species delimitations, evolutionary trajectories, and conservation strategies of the Galapagos giant tortoises. *Peer Community in Evolutionary Biology*, [100031]. <https://doi.org/10.24072/pci.evolbiol.100031>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.



A genomic perspective is needed for the re-evaluation of species boundaries, evolutionary trajectories, and conservation strategies of the Galápagos giant tortoises

Michael C Fontaine

Groningen Institute for Evolutionary Life Sciences (GELIFES). University of Groningen -- Groningen, The Netherlands

m.c.fontaine@rug.nl

doi: [10.24072/pci.evolbiol.100031](https://doi.org/10.24072/pci.evolbiol.100031)

Cite as: Fontaine MC. 2017. A genomic perspective is needed for the re-evaluation of species boundaries, evolutionary trajectories, and conservation strategies of the Galápagos giant tortoises. *Peer Community in Evolutionary Biology*, 100031. doi: [10.24072/pci.evolbiol.100031](https://doi.org/10.24072/pci.evolbiol.100031)

Open Access

A recommendation of

Loire E, Galtier N. 2017. **Lacking conservation genomics in the giant Galápagos tortoise.** *BiorXiv*, 101980, ver. 4 of 26th September 2017. doi: [10.1101/101980](https://doi.org/10.1101/101980)

Published: 26th September 2017

Copyright: This work is licensed under the Creative Commons Attribution-NoDerivatives 4.0 International License. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nd/4.0/>

Genome-wide data obtained from even a small number of individuals can provide unprecedented levels of detail about the evolutionary history of populations and species [1], determinants of genetic diversity [2], species boundaries and the process of speciation itself [3]. Loire and Galtier [4] present a clear example, using the emblematic Galápagos giant tortoise (*Chelonoidis nigra*), of how multi-species comparative population genomic approaches can provide valuable insights about population structure and species delimitation even when sample sizes are limited but the number of loci is large and distributed across the genome.

Galápagos giant tortoises are endemic to the Galápagos Islands and are currently recognized as an endangered, multi-species complex including both extant and extinct taxa. Taxonomic definitions are based on morphology, geographic isolation and population genetic evidence based on short DNA sequences of the mitochondrial genome (mtDNA) and/or a dozen or so nuclear microsatellite loci [5-8]. The species complex enjoys maximal protection. Population recoveries have been quite successful and spectacular conservation programs based on mitochondrial genes and microsatellites are ongoing. This includes for example individual translocations, breeding program, “hybrid” sterilization or removal, and resurrection of extinct lineages).

In 2013, Loire et al. [9] provided the first population genomic analyses based on genome scale data (~1000 coding loci derived from blood-transcriptomes) from five individuals, encompassing three putative “species”: *Chelonoidis becki*, *C. porteri* and *C. vandenburghi*. Their results raised doubts about the validity/accuracy of the currently accepted designations of “genetic distinctiveness”. However, the implications for conservation and management have remained unnoticed.

In 2017, Loire and Galtier [4] have re-appraised this issue using an original multi-species comparative population genomic analysis of their previous data set [9]. Based on a comparison of 53 animal species, they show that both the level of genome-wide neutral diversity (π_S) and level of population structure estimated using the inbreeding coefficient (F) are much lower than would be expected from a sample covering multiple species. The observed values are more comparable to those typically reported at the “among population” level within a single species such as human (*Homo sapiens*). The authors go to great length to assess the sensitivity of their method to detect population structure (or lack thereof) and show that their results are robust to potential issues, such as contamination and sequencing errors that can occur with Next Generation Sequencing techniques; and biases related to the small sample size and sub-sampling of individuals. They conclude that published mtDNA and microsatellite-based assessment of population structure and species designations may be biased towards over-splitting.

This manuscript is a very good read as it shows the potential of the now relatively affordable genome-wide data for helping to both resolve and clarify population and species boundaries, illuminate demographic trends, evolutionary trajectories of isolated groups, patterns of connectivity among them, and test for evidence of local adaptation and even reproductive isolation. The comprehensive information provided by genome-wide data can critically inform and assist the development of the best strategies to preserve endangered populations and species. Loire and Galtier [4] make a strong case for applying genomic data to the Galápagos giant tortoises, which is likely to redirect conservation efforts more effectively and at lower cost. The case of the Galápagos giant tortoises is certainly a very emblematic example, which will find an echo in many other endangered species conservation programs.

References

- [1] Li H and Durbin R. 2011. Inference of human population history from individual whole-genome sequences. *Nature*, 475: 493–496. doi: [10.1038/nature10231](https://doi.org/10.1038/nature10231)
- [2] Romiguier J, Gayral P, Ballenghien M, Bernard A, Cahais V, Chenuil A, Chiari Y, Dernat R, Duret L, Faivre N, Loire E, Lourenco JM, Nabholz B, Roux C, Tsagkogeorga G, Weber AA-T, Weinert LA, Belkhir K, Bierne N, Glémin S and Galtier N. 2014. Comparative population genomics in animals uncovers the determinants of genetic diversity. *Nature*, 515: 261–263. doi: [10.1038/nature13685](https://doi.org/10.1038/nature13685)
- [3] Roux C, Fraïsse C, Romiguier J, Anciaux Y, Galtier N and Bierne N. 2016. Shedding light on the grey zone of speciation along a continuum of genomic divergence. *PLoS Biology*, 14: e2000234. doi: [10.1371/journal.pbio.2000234](https://doi.org/10.1371/journal.pbio.2000234)

- [4] Loire E and Galtier N. 2017. Lacking conservation genomics in the giant Galápagos tortoise. bioRxiv 101980, ver. 4 of 26th September 2017. doi: [10.1101/101980](https://doi.org/10.1101/101980)
- [5] Beheregaray LB, Ciofi C, Caccone A, Gibbs JP and Powell JR. 2003. Genetic divergence, phylogeography and conservation units of giant tortoises from Santa Cruz and Pinzón, Galápagos Islands. *Conservation Genetics*, 4: 31–46. doi: [10.1023/A:1021864214375](https://doi.org/10.1023/A:1021864214375)
- [6] Ciofi C, Milinkovitch MC, Gibbs JP, Caccone A and Powell JR. 2002. Microsatellite analysis of genetic divergence among populations of giant Galápagos tortoises. *Molecular Ecology*, 11: 2265–2283. doi: [10.1046/j.1365-294X.2002.01617.x](https://doi.org/10.1046/j.1365-294X.2002.01617.x)
- [7] Garrick RC, Kajdacs B, Russello MA, Benavides E, Hyseni C, Gibbs JP, Tapia W and Caccone A. 2015. Naturally rare versus newly rare: demographic inferences on two timescales inform conservation of Galápagos giant tortoises. *Ecology and Evolution*, 5: 676–694. doi: [10.1002/ece3.1388](https://doi.org/10.1002/ece3.1388)
- [8] Poulakakis N, Edwards DL, Chiari Y, Garrick RC, Russello MA, Benavides E, Watkins-Colwell GJ, Glaberman S, Tapia W, Gibbs JP, Cayot LJ and Caccone A. 2015. Description of a new Galápagos giant tortoise species (*Chelonoidis*; Testudines: Testudinidae) from Cerro Fatal on Santa Cruz island. *PLoS ONE*, 10: e0138779. doi: [10.1371/journal.pone.0138779](https://doi.org/10.1371/journal.pone.0138779)
- [9] Loire E, Chiari Y, Bernard A, Cahais V, Romiguier J, Nabholz B, Lourenço JM and Galtier N. 2013. Population genomics of the endangered giant Galápagos tortoise. *Genome Biology*, 14: R136. doi: [10.1186/gb-2013-14-12-r136](https://doi.org/10.1186/gb-2013-14-12-r136)

Appendix

Reviews by four anonymous referees: <http://dx.doi.org/10.24072/pci.evolbiol.100031>